arXiv:0804.2736v1 [cond-mat.str-el] 17 Apr 2008

Non-analytic behavior of 2D itinerant ferromagnets.

Dmitry V. Efremov,[1] Joseph J. Betouras,[2] and Andrey Chubukov[3]

[1] Institute for Theoretical Physics, Technical University of Dresden, 01062 Dresden, Germany.
[2] Scottish Universities Physics Alliance, Department of Physics and Astronomy,
University of St. Andrews, St. Andrews KY16 9SS, Scotland, UK.
[3] Department of Physics, University of Wisconsin-Madison,
1150 University Ave.,Madison, WI 53706, USA.

(Dated: April 17, 2008)

We consider an ordered ferromagnet in the vicinity of a $T = 0$ transition into a paramagnet. We show that the free energy and the transverse and longitudinal static susceptibilities contain non-analyticities which destroy a continuous second-order transition. Depending on the parameters, the transition either becomes first-order, or occurs via an intermediate spiral phase.

PACS numbers:

Introduction. In recent years, there has been a strong interest to understand hidden features of a $T = 0$ ferromagnetic transition in itinerant fermionic systems. A much studied Hertz-Millis-Moriya (HMM) [1, 2, 3] ϕ^4 model of a ferromagnetic quantum criticality predicts that the transition should be continuous in all dimensions $D > 1$, with mean-field exponents, like a classical transition in $D > 4 = 1+z$, where $z = 3$ is the dynamical exponent. However, it was realized [4, 5, 6, 7] that the difference between quantum and classical cases is more than the change of the effective dimension – in the quantum case, the ϕ^4 and higher-order terms in the expansion in the order parameter field contain singular dynamic parts, which reflect the fact that fermions give rise to long-range, dynamical interaction between collective spin excitations in itinerant fermionic systems. These extra dynamic terms have been analyzed on the paramagnetic side of the transition, both in $D = 3$ (Ref. 5) and $D = 2$ (Ref. 8), and have been found to give rise to two competing effects: (i) the expansion of the free energy in the magnetic field is non-analytic, and the non-analytic term favors a preemptive first-order transition to a state with a finite magnetization, (ii) the static spin susceptibility $\chi(q)$ is non-analytic in q and becomes negative at some q_0, signaling another preemptive instability, this time towards a spiral phase. Which instability of a paramagnet comes first depends on the interplay between the prefactors for the analytic Δ^4 and q^2 terms, but in any event, a continuous second-order ferromagnetic transition point is internally unstable.

In this paper, we consider what happens when the system approaches a ferromagnetic quantum-critical point (QCP) from the ordered state. We show that the free energy of a quantum ferromagnet is non-analytic in the order parameter field Δ, and the non-analytic term favors a first-order transition into a paramagnet at some Δ_c. We analyze the static spin susceptibility, which in the ferromagnetic phase has non-equal longitudinal and transverse components, $\chi_\parallel(q)$ and $\chi_\perp(q)$, and show that both are non-analytic in momentum q and both become negative at a finite q inside the ferromagnetic case, when Δ becomes smaller that some critical $\Delta_{c\perp}$ and $\Delta_{c\parallel}$. The negative $\chi_{\parallel,\perp}(q)$ implies the development of the spiral order, either along the direction of the magnetization, or in a transverse direction. We relate $\Delta_c, \Delta_{c\perp}$ and $\Delta_{c\parallel}$ with the parameters of the model and argue that if $\Delta_c > \Delta_{c\perp}, \Delta_{c\parallel}$, the transition is first order, otherwise the ferromagnetic phase first becomes unstable against a spiral.

Our analysis is based on the Eliashberg-type consideration near a ferromagnetic QCP. Such approximation has been justified in [6, 7, 8] and we assume that it is valid. For definiteness, below we only consider the case $D = 2$, where the non-analyticities are stronger than in $D = 3$.

The point of departure of our analysis is the spin-fermion model near a QCP. It describes fermions interacting with their own collective excitations in the spin channel, described by spin variables $\mathbf{S} = c^\dagger_\alpha \sigma_{\alpha\beta} c_\beta$. The model does not assume a long-range order a 'priori, and is described by the Hamiltonian with three terms: $H_f = \sum_{k,\alpha=\uparrow,\downarrow} \varepsilon_{\mathbf{k}} c^\dagger_{\mathbf{k},\alpha} c_{\mathbf{k},\alpha}$, which describes low-energy fermions with the dispersion $\varepsilon_{\mathbf{k}} \approx \varepsilon_F + v_F(k - k_F)$, $H_s = \sum_q \chi_0^{-1}(q) \mathbf{S}_{\mathbf{q}} \mathbf{S}_{-\mathbf{q}}$, which describes collective bosonic excitations with a bare static propagator $\chi_0(q)$, and the spin-fermion interaction term $H_{int} = g\ (1/N) \sum_{\mathbf{k},\mathbf{q},\alpha,\alpha'} c^\dagger_{\mathbf{k},\alpha} \sigma_{\alpha,\alpha'} \mathbf{S}_{\mathbf{q}} c_{\mathbf{k}+\mathbf{q},\alpha'}$. Within this model, $\chi(q, \Omega_m) = \chi_0(q)/(1 - 2g^2\chi_0(q)\Pi(q, \Omega_m))$ where $\chi_0(q)$ is the static susceptibility of free fermions, and $\Pi(q, \Omega_m)$ is the polarization operator (here and below we set the Bohr magneton $\mu_B = 1$). Near a ferromagnetic transition, $\chi(q) = 2\nu(\varepsilon_F)/(\delta + (aq)^2)$, where $\nu(\varepsilon_F)$ is the density of states per particle at the Fermi surface ($\nu(\varepsilon_F) = m/(2\pi)$ for $\varepsilon_0(k) = k^2/(2m)$), $\delta = 1 - (2g\nu(\varepsilon_F))^2$, and the length a is proportional to the radius of the interaction [8]. The dynamic fermionic self-energy and the dynamic part of $\Pi(q, \Omega_m)$ are computed self-consistently within the model. The fully renormalized spin susceptibility in the paramagnetic phase is

$$\chi_{ij}(q,\Omega_m) = \delta_{ij}\ \frac{2\nu(\varepsilon_F)}{\delta + (aq)^2 + \bar{\Pi}(q,\Omega_m)}, \tag{1}$$

where $\bar{\Pi}(q,\Omega_m) = \Pi(q,\Omega_m) - \Pi(q,0)$. For free fermions $\bar{\Pi}(q,\Omega_m) = |\Omega_m|/\sqrt{(v_F q)^2 + \Omega_m^2}$.

The ferromagnetic transition occurs when $g = 1/(2\nu(\varepsilon_F))$. At larger g, δ becomes negative, and the system develops a ferromagnetic long-range order with the magnetization $M = \langle S^z\rangle = (N_\uparrow - N_\downarrow)$. We assume that such long-range order does exist and search for preemptive instabilities upon approaching QCP from the ferromagnetic side.

Mean-field analysis At the mean-field level, the interaction term reduces to $g\langle S_z\rangle \sum_{k,\alpha} \text{sign}\alpha\ c^\dagger_{\mathbf{k},\alpha} c_{\mathbf{k},\alpha}$, and the fermionic propagator becomes

$$G^{-1}_{\uparrow,\downarrow}(k,\omega_m) = i\omega_m - (\varepsilon_k - \mu) \pm \Delta_0, \tag{2}$$

where $\Delta_0 = g\langle S_z\rangle = g(N_\uparrow - N_\downarrow)$, and $\mu = \mu(\Delta_0)$ is the exact chemical potential (to order Δ_0^2, $\mu(\Delta_0) = \mu(0) - \nu'/(2\nu)\Delta_0^2$, where ν' is the derivative of the density of states at the Fermi surface). Because Δ_0 is finite, the longitudinal and transverse spin propagators become unequal already in the static limit. We have

$$\chi^{zz}(q) = \frac{\chi_0^{zz}}{1 - (g\chi_0^{zz})^2 + (aq)^2},$$
$$\chi^{xx}(q) = \frac{\chi_0^{xx}}{1 - (g\chi_0^{xx})^2 + (aq)^2}, \tag{3}$$

where

$$\chi_0^{zz} = -\int \frac{d\omega_m}{2\pi} \int \nu(\varepsilon) d\varepsilon \left(G^2_\uparrow(\varepsilon,\omega_m) + G^2_\downarrow(\varepsilon,\omega_m)\right),$$
$$\chi_0^{xx} = -2\int \frac{d\omega_m}{2\pi} \int \nu(\varepsilon) d\varepsilon G_\uparrow(\varepsilon,\omega_m) G_\downarrow(\varepsilon,\omega_m). \tag{4}$$

A simple calculation shows that $\chi^{xx}(0) = (N_\uparrow - N_\downarrow)/\Delta_0 = 1/g$, such that $\chi^{xx}(q)$ diverges at $q \to 0$, in agreement with the Goldstone theorem. For longitudinal susceptibility, $\chi^{zz}(q) = 2\nu(\varepsilon_F)/(\delta_F + (aq)^2)$, where $\delta_F = -2\delta = K\Delta_0^2$, and $K = -(2/3)\nu''/\nu$ (Ref.9). The theory is only valid when $K > 0$, otherwise the transition is first order by trivial reasons. By order of magnitude, $\delta_F \sim (\Delta_0/\varepsilon_F)^2$.

The dynamic terms $\bar{\Pi}^{xx}(q,\Omega_m)$ and $\bar{\Pi}^{zz}(q,\Omega_m)$ also differ at $\Delta_0 \neq 0$. Evaluating them using fermionic propagators from (2), we obtain

$$\bar{\Pi}^{zz}(q,\Omega_m) = \frac{|\Omega_m|}{\sqrt{(v_F q)^2 + \Omega_m^2}},$$
$$\bar{\Pi}^{xx}(q,i\Omega_m) = \frac{|\Omega_m|}{\sqrt{(v_F q)^2 + (\Omega_m + 2i\Delta_0)^2}}. \tag{5}$$

The mean-field dynamic spin susceptibilities in the ferromagnetic phase are then given by

$$\chi^{zz}(q,\Omega_m) = \frac{2\nu(\varepsilon_F)}{\delta_F + (aq)^2 + \bar{\Pi}^{zz}(q,\Omega_m)},$$
$$\chi^{xx}(q,i\Omega_m) = \frac{2\nu(\varepsilon_F)}{(aq)^2 + \bar{\Pi}^{xx}(q,\Omega_m)}. \tag{6}$$

Eliashberg theory Eqs. (2-6) constitute the mean-field description of the ferromagnetic phase. Within this description, the transition is continuous, i.e., the ferromagnetic phase is stable up to a point where $\Delta_0, \delta = 0$. The Eliashberg theory goes beyond this approximation – it self-consistently takes into account ω-dependent fermionic self-energy, but neglects $k-$dependent self-energy and vertex corrections. Vertex corrections generally are not small if the interaction involve small momentum transfers, and are necessary to satisfy Ward identities related to the conservation laws. However, the analysis of vertex corrections on the paramagnetic side have shown [7, 8] that they can be rigorously neglected in the calculations of the non-analytic terms in the free energy and spin susceptibilities, if the interaction is sufficiently long-ranged such that $ak_F > 1$, which we assume to hold [10].

The calculations proceed in three steps, like in the paramagnetic phase. First, we obtain self-consistent one-loop expressions for the fermionic self-energy and dynamic spin susceptibilities. Second, we use these one-loop expressions as inputs, obtain the free energy within Eliashberg theory, and show that it is non-analytic in Δ. Third, using the same inputs, we compute static spin susceptibilities at the two-loop level and find terms which are non-analytic in momentum. We argue that the non-analytic terms in the free energy favor a first-order transition, while the non-analytic terms in the susceptibilities favor an intermediate spiral phase.

Fermionic self-energy The one-loop fermionic self-energy in the ordered phase is given by

$$\Sigma_f(\omega_m) = \begin{cases} \lambda(\varepsilon_k)\ \omega_m & \text{for} \quad \omega_m \ll \omega_0/\lambda^3 \\ \omega_0^{1/3}\omega_m^{2/3} & \text{for} \quad \omega_m \gg \omega_0/\lambda^3 \end{cases}$$

where $\omega_0 = 3\sqrt{3}\varepsilon_F/(4(ak_F)^4)$, $\varepsilon_F = v_F k_F/2$, and $\lambda(\varepsilon_k)$ depends in non-singular way on the ratio of ε_k and Δ_0. For $\varepsilon_k = \mu$

$$\lambda = \frac{1}{4ak_F}\left(\frac{v_F}{a\Delta_0} + \frac{1}{\sqrt{\delta_F + (2a\Delta_0/v_F)^2}}\right). \tag{7}$$

The self-energy is linear in ω_m at the smallest frequencies, but crosses over to the quantum-critical, $\omega_m^{2/3}$ behavior at frequencies larger than $\omega_0/\lambda^3 \sim \Delta_0^3/(\varepsilon_F^2(ak_F))$. Such self-energy does not destroy the ferromagnetic order and preserves a Fermi surface, but it is larger than ω_m near QCP, and has non-Fermi liquid form in between

$\omega_0/\lambda^3 \ll \omega_0$ and ω_0. The non-Fermi liquid behavior in the ferromagnetic state has been discussed from a different point of view in Ref. [11].

The one-loop dynamic polarization operators $\bar{\Pi}_1^{xx}(q,\Omega_m)$ and $\bar{\Pi}_1^{zz}(q,\Omega_m)$, re-evaluated with dressed fermions, are given by rather complex expressions. Like in previous studies [7, 8], we found that, for the calculations of the non-analytic terms in the free energy and spin susceptibilities, we only need terms up to order $1/q^3$ in the $1/q$ expansion of $\bar{\Pi}(q,\Omega_m)$. Such terms are not affected by vertex corrections [7]. We obtained

$$\begin{aligned}\bar{\Pi}_1^{xx}(q,\omega) &= \frac{|\omega|}{v_F q}\left(1 - \frac{(\omega + c_\omega \Sigma_f(\omega) + 2i\Delta_0)^2}{2v_F^2 q^2}\right),\\ \bar{\Pi}_1^{zz}(q,\omega) &= \frac{|\omega|}{v_F q}\left(1 - \frac{(\omega + c_\omega \Sigma_f(\omega))^2}{2v_F^2 q^2}\right). \qquad (8)\end{aligned}$$

where c_ω interpolates between $c_{\omega=0} = 1$ and $c_\omega \approx 1.20$ for $\omega_0/\lambda^3 < \omega < \omega_0$.

The free energy The free energy per particle for the ferromagnetic spin-fermion model in the Eliashberg approximation is given by

$$\Xi = \Xi_0(\Delta_0) + \Xi^{zz} + 2\Xi^{xx},$$

where $\Xi_0(\Delta) = \frac{2m}{p_F^2}\left[-(1/4)\delta_F\Delta^2 + (K/8)\Delta^4 + ...\right]$ is analytic in Δ Minimizing Ξ_0 and expanding around the minimum, we obtain the equilibrium $\Delta = \Delta_0 = (\delta_F/K)^{1/2}$, and reproduce the mean-field expression for the static $\chi^{zz}(q \to 0)$. Further,

$$\begin{aligned}\Xi^{zz} &= \frac{V}{2N}\int \frac{d\Omega_m d^2q}{(2\pi)^3}\ln\frac{\chi^{zz}(0,0)}{\chi^{zz}(q,\Omega_m)},\\ \Xi^{xx} &= \frac{V}{2N}\int \frac{d\Omega_m d^2q}{(2\pi)^3}\ln\frac{\chi^{xx}(0,0)}{\chi^{xx}(q,\Omega_m)}, \qquad (9)\end{aligned}$$

where $\chi^{ii}(q,\Omega_m)$ include one-loop polarization operators $\bar{\Pi}_1^{ii}(q,\Omega_m)$. Both Ξ^{zz} and Ξ^{xx} contain analytic contributions which renormalize constants in $\Xi_0(\Delta)$. These renormalizations are small in $1/(ak_F)$ (Ref.[8]) and we neglect them. In addition, Ξ^{xx} contains the non-analytic term in Δ, which is our primary interest. Substituting χ^{xx} with the polarization operator from (8) into Eqn. (9), integrating over momentum, and neglecting regular terms, we obtain at QCP

$$\Xi^{xx} = \frac{\sqrt{2}}{2\pi c^{3/2}}\,\frac{\Delta^{7/2}}{\varepsilon_F^2\omega_0^{1/2}}Z, \qquad (10)$$

where Z is the universal (cutoff independent) part of the integral

$$Z = 2\int_0^A dyRe\left[(y^{2/3} - i)^2 \log{(y^{2/3} - i)^2}\right]. \qquad (11)$$

The evaluation of the integral yields $Z = -8\pi\sqrt{2}/35 \approx -1.02$. Combining non-analytic and analytic terms, we obtain in the immediate vicinity of the QCP

$$\Xi = \frac{1}{\varepsilon_F}\left[-\frac{\delta_F}{4}\Delta^2 + \frac{K}{8}\Delta^4 - \frac{\Delta^{7/2}}{E^{3/2}}\right], \qquad (12)$$

where $E = d\varepsilon_F/(ak_F)^{4/3}$, and $d \approx 3.50$. Apart for a small numerical difference in d, this expression coincides with the one obtained in [8], where QCP was approached from the paramagnetic side. There is, however, an important distinction between paramagnetic and ferromagnetic phases away from QCP. In the paramagnetic phase, the $\Delta^{7/2}$ dependence is replaced by Δ^3 when $\Delta/\varepsilon_F \sim \delta/(ak_F)^2 \leq \delta$. In the ferromagnetic phase, $\delta \to \delta_F$, which by itself scales as $(\Delta/\varepsilon_F)^2$. As a result, the non-analytic $\Delta^{7/2}$ dependence of Ξ survives in the wide range away from a QCP, as long as $\Delta/\varepsilon_F \leq \gamma$, where $\gamma = (ak_F)^2/(K\varepsilon_F^2)$.



FIG. 1: The free energy $\Xi(\Delta)$, Eq. (12), for different values of δ_F, going from positive $(\delta_F)_1$ to negative $(\delta_F)_{2,3,4}$. The first-order transition occurs at $(\delta_F)_3$.

The negative $\Delta^{7/2}$ term in the free energy shifts the equilibrium value Δ_0 such that it remains finite even on the paramagnetic side of the transition, when δ_F changes sign and becomes negative. For a generic δ_F, Δ_0 is the solution of $K\Delta_0^2 - 7(\Delta_0/E)^{3/2} = \delta_F$. One can easily verify (see Fig.1) that the free energy (12) describes a first-order transition to a paramagnet at $\delta_F = -(\Delta_0/E)^{3/2} < 0$ (i.e., $2\delta = -\delta_F > 0$). The value of the equilibrium Δ_0 at such transition is $\Delta_c = 36/(K^2E^3) \approx 0.84\gamma^2\varepsilon_F$. We also note that the stiffness – a prefactor for $(\Delta - \Delta_0)^2/2$ in the free energy – changes from δ_F to $\delta_F^{eff} = \delta_F + (7/4)(\Delta_0/E)^{3/2}$, and remains positive for all $\Delta_0 > \Delta_c$. At $\Delta_0 = \Delta_c$, $\delta_F^{eff} = (3/4)|\delta_F|$.

Static spin susceptibilities We next show that the static spin susceptibilities $\chi^{xx}(q,0)$ and $\chi^{zz}(q,0)$ also display non-analytic behavior, and that these non-analyticities compete with the one in the free energy and may give rise to pre-emptive spiral instabilities. The non-analytic term in $\chi(q,0)$ has been previously analyzed on the paramagnetic side [7, 8]. We performed the calculations in the ordered phase.

The non-analytic behavior of $\chi(q,0)$ originates from non-analytic $q-$dependencies of Π^{zz} and Π^{xx}, which ac-



FIG. 2: The scaling function $\sqrt{y}F^{xx}(y)$. It has a maximum value of ~ 0.24 at $y \approx 0.74$



FIG. 3: The three critical Δ_c as functions of the parameter $\gamma = (ak_F)^2/(K\varepsilon_F^2)$.

quire static parts at the two-loop order.

The computational steps are similar to those in Ref.[7] and we refrain from discussing them. The non-analytic contributions Π_2^{zz} and Π_2^{xx} come from the processes in which fermions and spin fluctuations are vibrating near a fermionic mass shell and are far away from a bosonic mass shell [6, 7] (the same processes lead to fermionic self-energy $\Sigma_f(\omega_m)$). We found that the non-analyticity comes from the exchange processes involving transverse spin fluctuations.

There are two types of non-analyticities in an ordered ferromagnet. First, there are corrections to static, uniform $\Pi^{zz}(0,0)$. They change δ_F into δ_F^{eff}, which is the same as the stiffness obtained by expanding the free energy. Second, there are non-analytic terms in the momentum expansion of the static $\Pi^{zz}(q,0)$ and $\Pi^{xx}(q,0)$ For the latter, we found

$$\begin{aligned}\Pi_2^{zz}(q,0) &= -a^2 q^{3/2} k_F^{1/2} F^{zz}(2\Delta_0/v_F q),\\ \Pi_2^{xx}(q,0) &= -a^2 q^{3/2} k_F^{1/2} F^{xx}(2\Delta_0/v_F q), \qquad (13)\end{aligned}$$

such that

$$\begin{aligned}\chi^{zz}(q,0) &= \frac{2\nu(\varepsilon_F)}{a^2}\ \frac{1}{\frac{\delta_F^{eff}}{a^2} + q^2 - q^{3/2}k_F^{1/2}F^{zz}\left(\frac{2\Delta_0}{v_F q}\right)},\\ \chi^{xx}(q,0) &= \frac{2\nu(\varepsilon_F)}{a^2}\ \frac{1}{q^2 - q^{3/2}k_F^{1/2}F^{xx}\left(\frac{2\Delta_0}{v_F q}\right)}. \qquad (14)\end{aligned}$$

For $\Delta_0 \ll v_F q$, $F^{zz}(0) = F^{xx}(0) \approx 0.25$, in agreement with [7]. For $v_F q \ll \Delta_0$ the two scaling functions differ. The scaling function $F^{zz}(y)$ remains close to $F^{zz}(0)$ as long as the argument $y < \varepsilon_F/\Delta_0$. The function $F^{xx}(y)$ crosses over to $F^{xx}(y \gg 1) \approx 0.15/\sqrt{y}$ such that $\Pi_2^{xx}(q,0)$ becomes analytic, $\Pi_2^{xx}(q,0) \propto q^2$. The function $\sqrt{y}F^{xx}(y)$ is plotted in Fig. 2 Analyzing Eq. (14), we find that both susceptibilities become negative at a finite q when Δ_0 reduces below some critical value. The transverse susceptibility becomes negative at $\Delta_0 = \Delta_{c,\perp} = \varepsilon_F(\sqrt{y}F(y))^2_{max} = 0.06\varepsilon_F$, whereas longitudinal susceptibility becomes negative at $\Delta_0 = \Delta_{c,\|}$, which is the solution of

$$\frac{\Delta_{c,\|}}{\varepsilon_F}\left(1 - \frac{21}{4}\frac{\gamma}{d^{3/2}}\sqrt{\frac{\varepsilon_F}{\Delta_{c,\|}}}\right)^{1/2} = 0.02\sqrt{\gamma}.$$

Whether the transition is the first order or involves an intermediate spiral phase depends on which of Δ_c, $\Delta_{c,\|}$ and $\Delta_{c,\perp}$ is the largest. All three critical Δ scale with ε_F, but they depend differently on the parameter γ. In Fig.3 we plotted the three critical Δ vs γ. We see that for $\gamma < 0.26$, $\Delta_{c,\perp} > \Delta_{c,\|}$, Δ_c, and the system first develops a transverse spiral order, while for $\gamma > 0.26$, $\Delta_c > \Delta_{c,\|}$, $\Delta_{c,\perp}$, and the transition is first order. This qualitatively agrees with the analysis on the paramagnetic side [8].

In summary, we analyzed the non-analytic terms in the free energy and in static susceptibilities in an ordered itinerant ferromagnet. We found that, because of these non-analyticities, the transition to a paramagnet is either first order, or involves an intermediate spiral phase.

We thank A. Andreev, P. Fulde, A. Green, A. Huxley, D. Maslov, A. Rosch, and Yu. Ovchinnikov for useful discussions. The research has been supported by NSF DMR 0604406 (A. Ch.) and in part by NSF PHY05-51164 (J.B.). J.B. would like to thank KITP, Santa Barbara for hospitality during the completion of this work.

[1] J. Hertz, Phys. Rev. B **14**, 1165 (1976).
[2] T. Moriya,"Spin fluctuations in itinerant electron magnetism", eds Springer-Verlag, 1985.
[3] A. J. Millis, Phys. Rev. B **48**, 7183 (1993).
[4] D. Belitz, T. R. Kirkpatrick, and T. Vojta Rev. Mod. Phys. **77**, 579 (2005)
[5] D. Belitz, T. R. Kirkpatrick, and T. Vojta, Phys. Re v. B **55**, 9452 (1997).
[6] A. V. Chubukov, C. Pepin, and J. Rech, Phys. Rev. Lett. **92**, 147003 (2004).
[7] J. Rech, C. Pepin, and A. V. Chubukov, Phys. Rev. B **74**, 195126 (2006).
[8] D. L. Maslov, A. V. Chubukov, and R. Saha, Phys. Rev. B. **74**, 220402(R) (2006).

[9] A somewhat different result for K is obtained if one uses RPA for the Hubbard model. In this case, charge component of the interaction also contributes to K, and $K = (\nu'/\nu)^2 - (2/3)\nu''/\nu$.

[10] An alternative way to justify Eliashberg theory even if $ak_F \leq 1$ is to extend the model to $M \gg 1$ spieces of fermions, which separately interact with a collective boson. Vertex corrections are then small in $1/M$.

[11] V.P. Mineev, Phys. Rev. B**72**, 144418 (2005); D. Belitz, T. R. Kirkpatrick, and A. Rosch, Phys. Rev. B **74**, 024409 (2006).